PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: WO 96/33735 (11) International Publication Number: A61K 39/00, C07K 16/18, C12N 5/16, A1 (43) International Publication Date: 31 October 1996 (31.10.96) (21) International Application Number: PCT/US96/05928 (81) Designated States: AU, CA, HU, JP, KR, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, (22) International Filing Date: 29 April 1996 (29.04.96) LU, MC, NL, PT, SE). (30) Priority Data: **Published** 430,938 27 April 1995 (27.04.95) US With international search report. (71) Applicant: CELL GENESYS, INC. [US/US]; 344 Lakeside Drive, Foster City, CA 94404 (US). (72) Inventors: KUCHERLAPATI, Raju; 8 Gracie Lane, Darien, CT 06820 (US). JAKOBOVITS, Aya: 2021 Monterey Avenue, Menlo Park, CA 94025 (US). KLAPHOLZ. Sue: 76 Peter Courts Circle, Stanford, CA 94305 (US). BRENNER, Daniel, G.; 86 Central Avenue, Redwood City, CA 94601 (US). CAPON, Daniel, J.; 90 Woodridge Road, Hillsborough, CA 94010 (US). (74) Agents: HALLUIN, Albert, P. et al.; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).

(54) Title: HUMAN ANTIBODIES DERIVED FROM IMMUNIZED XENOMICE

(57) Abstract

Fully human antibodies against a specific antigen can be prepared by administering the antigen to a transgenic animal which has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled. Various subsequent manipulations can be performed to obtain either antibodies per se or analogs thereof.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Beigium	HU	Hungary	NO	Norway
BF	Burkma Faso	1E	Ireland	NZ	New Zealand
BG	Bulgana	IT	Italy	PL	Poland
BJ	Benin	JР	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CC	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'ivoire	u	Liechtensiein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	Chma	LR	Liberta	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TC	Togo
DE	Germany	LV	Laivia	TJ	Tajikistan
DK	Denmark	MC	Monaco	ıπ	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain _	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UŽ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Vici Nam

HUMAN ANTIBODIES DERIVED FROM IMMUNIZED XENOMICE

Technical Field

5

10

15

20

25

30

35

The invention relates to the field of immunology, and in particular to the production of antibodies. More specifically, it concerns producing such antibodies by a process which includes the step of immunizing a transgenic animal with an antigen to which antibodies are desired. The transgenic animal has been modified so as to produce human, as opposed to endogenous, antibodies.

Background Art

PCT application WO 94/02602, published 3 February 1994 and incorporated herein by reference, describes in detail the production of transgenic nonhuman animals which are modified so as to produce fully human antibodies rather than endogenous antibodies in response to antigenic challenge. Briefly, the endogenous loci encoding the heavy and light immunoglobulin chains are incapacitated in the transgenic hosts and loci encoding human heavy and light chain proteins are inserted into the genome. In general, the animal which provides all the desired modifications is obtained by cross breeding intermediate animals containing fewer than the full complement of modifications. preferred embodiment of nonhuman animal described in the specification is a mouse. Thus, mice, specifically, are described which, when administered immunogens, produce antibodies with human variable regions, including fully human antibodies, rather than murine antibodies that are immunospecific for these antigens.

The availability of such transgenic animals makes possible new approaches to the production of fully human antibodies. Antibodies with various immunospecificities are desirable for therapeutic and diagnostic use. Those antibodies intended for human therapeutic and in vivo diagnostic use, in particular, have been problematic because prior art sources for such antibodies resulted in immunoglobulins bearing the characteristic structures of

antibodies produced by nonhuman hosts. Such antibodies tend to be immunogenic when used in humans.

The availability of the nonhuman, immunogen responsive transgenic animals described in the above-referenced WO 94/02602 make possible convenient production of human antibodies without the necessity of employing human hosts.

Disclosure of the Invention

5

10

15

20

25

30

35

The invention is directed to methods to produce human antibodies by a process wherein at least one step of the process includes immunizing a transgenic nonhuman animal with the desired antigen. The modified animal fails to produce endogenous antibodies, but instead produces B-cells which secrete fully human immunoglobulins. The antibodies produced can be obtained from the animal directly or from immortalized B-cells derived from the animal. Alternatively, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly or modified to obtain analogs of antibodies such as, for example, single chain F, molecules.

Thus, in one aspect, the invention is directed to a method to produce a fully human immunoglobulin to a specific antigen or to produce an analog of said immunoglobulin by a process which comprises immunizing a nonhuman animal with the antigen under conditions that stimulate an immune response. The nonhuman animal is characterized by being substantially incapable of producing endogenous heavy or light immunoglobulin chain, but capable of producing immunoglobulins with both human variable and constant In the resulting immune response, the animal produces B cells which secrete immunoglobulins that are fully human and specific for the antigen. The human immunoglobulin of desired specificity can be directly recovered from the animal, for example, from the serum, or primary B cells can be obtained from the animal and immortalized. immortalized B cells can be used directly as the source of human antibodies or, alternatively, the genes encoding the

antibodies can be prepared from the immortalized B cells or from primary B cells of the blood or lymphoid tissue (spleen, tonsils, lymph nodes, bone marrow) of the immunized animal and expressed in recombinant hosts, with or without modifications, to produce the immunoglobulin or its analogs. In addition, the genes encoding the repertoire of

In addition, the genes encoding the repertoire of immunoglobulins produced by the immunized animal can be used to generate a library of immunoglobulins to permit screening for those variable regions which provide the desired

affinity. Clones from the library which have the desired characteristics can then be used as a source of nucleotide sequences encoding the desired variable regions for further manipulation to generate antibodies or analogs with these characteristics using standard recombinant techniques.

In another aspect, the invention relates to an immortalized nonhuman B cell line derived from the above described animal. In still another aspect, the invention is directed to a recombinant host cell which is modified to contain the gene encoding either the human immunoglobulin with the desired specificity, or an analog thereof which exhibits the same specificity.

In still other aspects, the invention is directed to antibodies or antibody analogs prepared by the above-described methods and to recombinant materials for their production.

In still other aspects, the invention is directed to antibodies which are immunospecific with respect to particular antigens set forth herein and to analogs which are similarly immunospecific, as well as to the recombinant materials useful to production of these antibodies.

Brief Description of the Drawings

Figure 1 is a schematic of the construction of the yH1C human heavy chain YAC.

Figure 2 is a schematic of the construction of the yK2 human kappa light chain YAC.

Figure 3 shows the serum titers of anti-IL-6 antibodies from a XenoMouse immunized with human IL-6 and

5

15

20

25

which antibodies contain human κ light chains and/or human μ heavy chains.

Figure 4 show the serum titers of anti-TNF α antibodies from a XenoMouse^M immunized with human TNF- α and which antibodies contain human κ light chains and/or human μ heavy chains.

Figure 5 shows serum titers of anti-CD4 antibodies from a XenoMouse immunized with human CD4 and which antibodies contain human κ light chains and/or human μ heavy chains.

Figure 6 shows the serum titers of a XenoMouse immunized with 300.19 cells expressing L-selectin at their surface. In the ELISA assay used, these antibodies are detectable if they carry human μ constant region heavy chains.

Figure 7 shows the serum titers of a XenoMouse^{∞} immunized with 300.19 cells expressing L-selectin at their surface. In the ELISA assay used, these antibodies are detectable only if they carry human κ light chains.

Figure 8 shows a FACS Analysis of human neutrophils incubated with serum from a XenoMouse immunized with human L-selectin and labeled with an antibody immunoreactive with human light chain κ region.

Figure 9 shows a diagram of a plasmid used to transfect mammalian cells to effect the production of the human protein gp39.

Figure 10 represents the serum titration curve of mice immunized with CHO cells expressing human gp39. The antibodies detected in this ELISA must be immunoreactive with gp39 and contain human heavy chain μ constant regions of human κ light chains.

Figure 11 is a titration curve with respect to monoclonal antibodies secreted by the hybridoma clone D5.1. This clone is obtained from a XenoMouse immunized with tetanus toxin C (TTC) and contains human κ light chain and human μ constant region in the heavy chain.

5

10

15

20

25

30

Figure 12 DNA sequence of the heavy chain of antitetanus toxin monoclonal antibody D5.1.4 (a subclone of D5.1). Mutations form germline are boxed.

Figure 13 DNA sequence of the kappa light chain of anti-tetanus toxin monoclonal antibody D5.1.4. Mutations form germline are boxed.

Figure 14 shows the serum titers of anti-IL-8 antibodies of XenoMouse^m immunized with human IL-8 and which antibodies contain human κ light chains and/or human μ heavy chains.

Figure 15 Inhibition of IL-8 binding to human neutrophils by monoclonal anti-human-IL-8 antibodies.

Figure 16 (A-H) DNA sequences of the heavy chain and kappa light chain of the anti-IL-8 antibodies D1.1 (16A-15 B), K2.2 (16C-D), K4.2 (16E-F), and K4.3 (16G-H).

Modes of Carrying Out the Invention

In general, the methods of the invention include administering an antigen for which human forms of immunospecific reagents are desired to a transgenic nonhuman 20 animal which has been modified genetically so as to be capable of producing human, but not endogenous, antibodies. Typically, the animal has been modified to disable the endogenous heavy and/or kappa light chain loci in its genome, so that these endogenous loci are incapable of the 25 rearrangement required to generate genes encoding immunoglobulins in response to an antigen. In addition, the animal will have been provided, stably, in its genome, at least one human heavy chain locus and at least one human light chain locus so that in response to an administered 30 antigen, the human loci can rearrange to provide genes encoding human variable regions immunospecific for the antigen.

The details for constructing such an animal useful in the method of the invention are provided in the PCT application WO 94/02602 referenced above. Examples of YACs for the present invention can be found in, for example, Green et al. Nature Genetics 7:13-21 (1994). In a preferred

5

embodiment of the XenoMouse™, the human heavy chain YAC, yH1C (1020 kb), and human light chain YAC, yK2 (880 kb) are used. yH1C is comprised of 870 kb of the human variable region, the entire D and J_u region, human μ , δ , and γ 2 constant regions and the mouse 3' enhancer. yK2 is comprised of 650 kb of the human kappa chain proximal variable region (Vx), the entire J_K region, and C_K with its flanking sequences that contain the Kappa deleting element (xde). Both YACs also contain a human HPRT selectable marker on their YAC vector arm. Construction of yH1C and yK2 was accomplished by methods well known in the art. In brief, YAC clones bearing segments of the human immunoglobulin loci were identified by screening a YAC library (Calbertsen et al, PNAS 87:4256 (1990)) Overlapping clones were joined by recombination using standard techniques (Mendez et al. Genomics 26:294-307 (1995)). Details of the schemes for assembling yH1C and yK2 are shown in Figure 1 and Figure 2 respectively.

yK2 was constructed from the clones A80-C7, A210-F10 and A203-C6 from the Olson library, disclosed in, for example, Burke et al., Science 236:806-812 (1987), Brownstein et al., Science 244:1348-1351 (1989), and Burke et al., Methods in Enzymology 194:251-270 (1991).

For production of the desired antibodies, the first step is administration of the antigen. Techniques for such administration are conventional and involve suitable immunization protocols and formulations which will depend on the nature of the antigen per se. It may be necessary to provide the antigen with a carrier to enhance its immunogenicity and/or to include formulations which contain adjuvants and/or to administer multiple injections and/or to vary the route of the immunization, and the like. Such techniques are standard and optimization of them will depend on the characteristics of the particular antigen for which immunospecific reagents are desired.

As used herein, the term "immunospecific reagents" includes immunoglobulins and their analogs. The term "analogs" has a specific meaning in this context. It refers to moieties that contain the fully human portions of the

5

10

15

20

25

30

immunoglobulin which account for its immunospecificity. In particular, complementarity determining regions (CDRs) are required, along with sufficient portions of the framework (Frs) to result in the appropriate three dimensional conformation. Typical immunospecific analogs of antibodies include F(ab")₂, Fab', and Fab regions. Modified forms of the variable regions to obtain, for example, single chain F, analogs with the appropriate immunospecificity are known. A review of such F, construction is found, for example, in Huston et al., Methods in Enzymology 203:46-63 (1991). The construction of antibody analogs with multiple immunospecificities is also possible by coupling the variable regions from one antibody to those of second antibody.

The variable regions with fully human characteristics can also be coupled to a variety of 15 additional substances which can provide toxicity, biological functionality, alternative binding specificities and the The moieties including the fully human variable regions produced by the methods of the invention include single-chain fusion proteins, molecules coupled by covalent 20 methods other than those involving peptide linkages, and aggregated molecules. Examples of analogs which include variable regions coupled to additional molecules covalently or noncovalently include those in the following nonlimiting illustrative list. Traunecker, A. et al. Int. J. Cancer Supp 25 (1992) Supp 7:51-52 describe the bispecific reagent janusin in which the F, region directed to CD3 is coupled to soluble CD4 or to other ligands such as OVCA and IL-7. Similarly, the fully human variable regions produced by the method of the invention can be constructed into $F_{\rm v}$ molecules and coupled 30 to alternative ligands such as those illustrated in the cited article. Higgins, P.J. et al J.Infect Disease (1992) 166:198-202 described a heteroconjugate antibody composed of OKT3 cross-linked to an antibody directed to a specific sequence in the V3 region of GP120. 35 Such heteroconjugate antibodies can also be constructed using at least the human variable regions contained in the immunoglobulins produced by the invention methods. Additional examples of bispecific

5

antibodies include those described by Fanger, M.W. et al.

Cancer Treat Res (1993) 68:181-194 and by Fanger, M.W. et al.

Crit Rev Immunol (1992) 12:101-124. Conjugates that are immunotoxins including conventional antibodies have been widely described in the art. The toxins may be coupled to the antibodies by conventional coupling techniques or immunotoxins containing protein toxin portions can be produced as fusion proteins. The analogs of the present invention can be used in a corresponding way to obtain such immunotoxins. Illustrative of such immunotoxins are those described by Byers, B.S. et al. Seminars Cell Biol (1991) 2:59-70 and by Fanger, M.W. et al. Immunol Today (1991) 12:51-54.

It will also be noted that some of the immunoglobulins and analogs of the invention will have agonist activity with respect to antigens for which they are immunospecific in the cases wherein the antigens perform signal transducing functions. Thus, a subset of antibodies or analogs prepared according to the methods of the invention which are immunospecific for, for example, a cell surface receptor, will be capable of eliciting a response from cells bearing this receptor corresponding to that elicited by the native ligand. Furthermore, antibodies or analogs which are immunospecific for substances mimicking transition states of chemical reactions will have catalytic activity. Hence, a subset of the antibodies and analogs of the invention will function as catalytic antibodies.

In short, the genes encoding the immunoglobulins produced by the transgenic animals of the invention can be retrieved and the nucleotide sequences encoding the fully human variable region can be manipulated according to known techniques to provide a variety of analogs such as those described above. In addition, the immunoglobulins themselves containing the human variable regions can be modified using standard coupling techniques to provide conjugates retaining immunospecific regions.

Thus, immunoglobulin "analogs" refers to the moieties which contain those portions of the antibodies of

5

10

15

20

25

30

the invention which retain their human characteristics and their immunospecificity. These will retain sufficient human variable regions to provide the desired specificity.

As stated above, all of the methods of the invention include administering the appropriate antigen to the transgenic animal. The recovery or production of the antibodies themselves can be achieved in various ways.

First, and most straightforward, the polyclonal antibodies produced by the animal and secreted into the bloodstream can be recovered using known techniques. Purified forms of these antibodies can, of course, be readily prepared by standard purification techniques, preferably including affinity chromatography with Protein A, anti-immunoglobulin, or the antigen itself. In any case, in order to monitor the success of immunization, the antibody levels with respect to the antigen in serum will be monitored using standard techniques such as ELISA, RIA and the like.

For some applications only the variable regions of the antibodies are required. Treating the polyclonal antiserum with suitable reagents so as to generate Fab', Fab, or F(ab"), portions results in compositions retaining fully human characteristics. Such fragments are sufficient for use, for example, in immunodiagnostic procedures involving coupling the immunospecific portions of immunoglobulins to detecting reagents such as radioisotopes.

Alternatively, immunoglobulins and analogs with desired characteristics can be generated from immortalized B cells derived from the transgenic animals used in the method of the invention or from the rearranged genes provided by these animals in response to immunization.

Thus, as an alternative to harvesting the antibodies directly from the animal, the B cells can be obtained, typically from the spleen, but also, if desired, from the peripheral blood lymphocytes or lymph nodes and immortalized using any of a variety of techniques, most commonly using the fusion methods described by Kohler and Milstein Nature 245:495 (1975) The resulting hybridomas (or otherwise immortalized B cells) can then be cultured as

5

10

15

20

25

30

single colonies and screened for secretion of antibodies of the desired specificity. As described above, the screen can also include a confirmation of the fully human character of the antibody. For example, as described in the examples below, a sandwich ELISA wherein the monoclonal in the hybridoma supernatant is bound both to antigen and to an antihuman constant region can be employed. After the appropriate hybridomas are selected, the desired antibodies can be recovered, again using conventional techniques. can be prepared in quantity by culturing the immortalized B cells using conventional methods, either in vitro or in vivo to produce ascites fluid. Purification of the resulting monoclonal antibody preparations is less burdensome that in the case of serum since each immortalized colony will secrete only a single type of antibody. In any event, standard purification techniques to isolate the antibody from other proteins in the culture medium can be employed.

As an alternative to obtaining human immunoglobulins directly from the culture of immortalized B cells derived from the animal, the immortalized cells can be used as a source of rearranged heavy chain and light chain loci for subsequent expression and/or genetic manipulation. Rearranged antibody genes can be reverse transcribed from appropriate mRNAs to produce cDNA. If desired, the heavy chain constant region can be exchanged for that of a different isotype or eliminated altogether. regions can be linked to encode single chain F, regions. Multiple F. regions can be linked to confer binding ability to more than one target or chimeric heavy and light chain combinations can be employed. Once the genetic material is available, design of analogs as described above which retain both their ability to bind the desired target, and their human characteristics, is straightforward.

Once the appropriate genetic material is obtained and, if desired, modified to encode an analog, the coding sequences, including those that encode, at a minimum, the variable regions of the human heavy and light chain, can be inserted into expression systems contained on vectors which

5

10

15

20

25

30

can be transfected into standard recombinant host cells. As described below, a variety of such host cells may be used; for efficient processing, however, mammalian cells are preferred. Typical mammalian cell lines useful for this purpose include CHO cells, 293 cells, or NSO cells.

The production of the antibody or analog is then undertaken by culturing the modified recombinant host under culture conditions appropriate for the growth of the host cells and the expression of the coding sequences. The antibodies are then recovered from the culture. The expression systems are preferably designed to include signal peptides so that the resulting antibodies are secreted into the medium; however, intracellular production is also possible.

15 In addition to deliberate design of modified forms of the immunoglobulin genes to produce analogs, advantage can be taken of phage display techniques to provide libraries containing a repertoire of antibodies with varying affinities for the desired antigen. For production of such repertoires, it is unnecessary to immortalize the B cells from the 20 immunized animal; rather, the primary B cells can be used directly as a source of DNA. The mixture of cDNAs obtained from B cells, e.g., derived from spleens, is used to prepare an expression library, for example, a phage display library transfected into E. coli. The resulting cells are tested for 25 immunoreactivity to the desired antigen. Techniques for the identification of high affinity human antibodies from such libraries are described by Griffiths, A.D., et al., EMBO J (1994) 13:3245-3260; by Nissim, A., et al. ibid, 692-698, and by Griffiths, A.D., et al., ibid, 12:725-734. Ultimately, 30 clones from the library are identified which produce binding affinities of a desired magnitude for the antigen, and the DNA encoding the product responsible for such binding is recovered and manipulated for standard recombinant expression. Phage display libraries may also be constructed 35 using previously manipulated nucleotide sequences and screened in similar fashion. In general, the cDNAs encoding

5

heavy and light chain are independently supplied or are linked to form F, analogs for production in the phage library.

The phage library is then screened for the antibodies with highest affinity for the antigen and the genetic material recovered from the appropriate clone. Further rounds of screening can increase the affinity of the original antibody isolated. The manipulations described above for recombinant production of the antibody or modification to form a desired analog can then be employed.

10 Combination of phage display technology with the XenoMouse™ offers a significant advantage over previous applications of phage display. Typically, to generate a highly human antibody by phage display, a combinatorial antibody library is prepared either from human bone marrow or 15 from peripheral blood lymphocytes as described by Burton, D.R., et al., Proc. Natl. Acad. Sci. USA (1991) 88:10134-10137. Using this approach, it has been possible to isolate high affinity antibodies to human pathogens from infected individuals, i.e. from individuals who have been "immunized" 20 as described in Burton, D.R., et al., Proc. Natl. Acad. Sci. <u>USA</u> (1991) <u>88</u>:10134-10137, Zebedee, S.L., et al. <u>Proc. Natl.</u> Acad. Sci. USA (1992) 89:3175-3179, and Barbas III, C.F., et al., Proc. Natl. Acad. Sci. USA (1991) 89:10164-20168. However, to generate antibodies reactive with human antigens, it has been necessary to generate synthetic libraries (Barbas 25 III C.F., et al., Proc. Natl. Acad. Sci. USA (1991) 89:4457-4461, Crameri, A. et al., BioTechniques (1995) 88:194-196) or to prepare libraries from either autoimmune patients (Rapoport, B., et al., <u>Immunol. Today</u> (1995) 16:43-49, 30 Portolano, S., et al., <u>J. Immunol.</u> (1993) <u>151</u>:2839-2851, and Vogel, M., et al., Eur J. Immunol. (1994) 24:1200-1207) or normal individuals, i.e. naive libraries (Griffiths, A.D., et al., <u>EMBO J.</u> (1994) <u>13</u>:3245-3260, Griffiths, A.D., et al., EMBO J. (1993) 12:725-734, Persson, M.A.A., et al., Proc. 35 Natl. Acad. Sci. USA (1991) 88:2432-2436, Griffiths, A.D., <u>Curr. Opin. Immunol.</u> (1993) <u>5</u>:263-267, Hoogenboom, H.R., et al., J. Mol. Biol. (1992) 227:381-388, Lerner, R.A., et al.,

Science (1992) 258:1313-1314, and Nissim A., et al., EMBO J.

(1994) 13:692-698. Typically, high affinity antibodies to human proteins have proven very difficult to isolate in this way. As is well known, affinity maturation requires somatic mutation and somatic mutation, in turn, is antigen driven.

- In the XenoMouse, repeated immunization with human proteins will lead to somatic mutation and, consequently, high affinity antibodies. The genes encoding these antibodies can be readily amplified by PCR as described in Marks, J.D., et al., J. Mol. Biol. (1991) 581-596 and immunospecific
- antibodies isolated by standard panning techniques, Winter, G., et al., Annu. Rev. Immunol. (1994) 12:433-55 and Barbas III, C.F., et al., Proc. Natl. Acad. Sci. USA (1991) 88:7978-7982.

As above, the modified or unmodified rearranged

loci are manipulated using standard recombinant techniques by
constructing expression systems operable in a desired host
cell, such as, typically, a Chinese hamster ovary cell, and
the desired immunoglobulin or analog is produced using
standard recombinant expression techniques, and recovered and
purified using conventional methods.

The application of the foregoing processes to antibody production has enabled the preparation of human immunospecific reagents with respect to antigens for which human antibodies have not heretofore been available. The immunoglobulins that result from the above-described methods and the analogs made possible thereby provide novel compositions for use in analysis, diagnosis, research, and therapy. The particular use will, of course, depend on the immunoglobulin or analog prepared. In general, the compositions of the invention will have utilities similar to those ascribable to nonhuman antibodies directed against the same antigen. Such utilities include, for example, use as affinity ligands for purification, as reagents in immunoassays, as components of immunoconjugates, and as therapeutic agents for appropriate indications.

Particularly in the case of therapeutic agents or diagnostic agents for use in *vivo*, it is highly advantageous to employ antibodies or their analogs with fully human

25

30

characteristics. These reagents avoid the undesired immune responses engendered by antibodies or analogs which have characteristics marking them as originating from nonhuman species. Other attempts to "humanize" antibodies do not result in reagents with fully human characteristics. For example, chimeric antibodies with murine variable regions and human constant regions are easily prepared, but, of course, retain murine characteristics in the variable regions. Even the much more difficult procedure of "humanizing" the variable regions by manipulating the genes encoding the amino acid sequences that form the framework regions does not provide the desired result since the CDRs, typically of nonhuman origin, cannot be manipulated without destroying immunospecificity.

Thus, the methods of the present invention provide, for the first time, immunoglobulins that are fully human or analogs which contain immunospecific regions with fully human characteristics.

There are large numbers of antigens for which human antibodies and their human analogs would be made available by the methods of the invention. These include, but are not limited to, the following nonlimiting set:

leukocyte markers, such as CD2, CD3, CD4, CD5, CD6, CD7, CD8, CD11a,b,c, CD13, CD14, CD18, CD19, CD20, CD22, CD23, CD27 and its ligand, CD28 and its ligands B7.1, B7.2, B7.3, CD29 and its ligand, CD30 and its ligand, CD40 and its ligand gp39, CD44, CD45 and isoforms, Cdw52 (Campath antigen), CD56, CD58, CD69, CD72, CTLA-4, LFA-1 and TCR

histocompatibility antigens, such as MHC class I or II, the Lewis Y antigens, Slex, Sley, Slea, and Selb;

adhesion molecules, including the integrins, such as VLA-1, VLA-2, VLA-3, VLA-4, VLA-5, VLA-6, LFA-1, Mac-1, $\alpha V\beta 3$, and p150,95; and

5

10

15

20

25

the selectins, such as L-selectin, E-selectin, and P-selectin and their counterreceptors VCAM-1, ICAM-1, ICAM-2, and LFA-3;

interleukins, such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, and IL-15;

interleukin receptors, such as IL-1R, IL-2R, IL-3R, ID-4R, IL-5R, IL-6R, IL-7R, IL-8R, IL-9R, IL-10R, IL-11R, IL-12R, IL-13R, IL-14R and IL-15R;

chemokines, such as PF4, RANTES, MIP1 α , MCP1, IP-10, ENA-78, NAP-2, Gro α , Gro β , and IL-8;

15

growth factors, such as TNFalpha, TGFbeta, TSH, VEGF/VPF, PTHrP, EGF family, FGF, PDGF family, endothelin, Fibrosin (F_sF_{-1}) , Laminin, and gastrin releasing peptide (GRP);

20

growth factor receptors, such as TNFalphaR, RGFbetaR, TSHR, VEGFR/VPFR, FGFR, EGFR, PTHrPR, PDGFR family, EPO-R, GCSF-R and other hematopoietic receptors;

interferon receptors, such as IFN α R, IFN β R, and IFN $_{\gamma}$ R;

Igs and their receptors, such as IGE, FceRI, and
FceRII;

30

tumor antigens, such as her2-neu, mucin, CEA and endosialin;

allergens, such as house dust mite antigen, lol pl (grass) antigens, and urushiol;

viral proteins, such as CMV glycoproteins B, H, and gCIII, HIV-1 envelope glycoproteins, RSV envelope

glycoproteins, HSV envelope glycoproteins, EBV envelope glycoproteins, VZV, envelope glycoproteins, HPV envelope glycoproteins, Hepatitis family surface antigens;

toxins, such as pseudomonas endotoxin and osteopontin/uropontin, snake venom, spider venom, and bee venom:

blood factors, such as complement C3b, complement

C5a, complement C5b-9, Rh factor, fibrinogen, fibrin, and

myelin associated growth inhibitor;

enzymes, such as cholesterol ester transfer protein, membrane bound matrix metalloproteases, and glutamic acid decarboxylase (GAD); and

miscellaneous antigens including ganglioside GD3, ganglioside GM2, LMP1, LMP2, eosinophil major basic protein, PTHrp, eosinophil cationic protein, pANCA, Amadori protein, Type IV collagen, glycated lipids, v-interferon, A7, P-glycoprotein and Fas (AFO-1) and oxidized-LDL.

Particularly preferred immunoglobulins and analogs are those immunospecific with respect to human IL-6, human IL-8, human TNFα, human CD4, human L-selectin, human PTHrp and human gp39. Antibodies and analogs immunoreactive with human TNF α and human IL-6 are useful in treating cachexia and septic shock as well as autoimmune disease. Antibodies and analogs immunoreactive with GP39 or with L-selectin are also effective in treating or preventing autoimmune disease. addition, anti-gp39 is helpful in treating graft versus host disease, in preventing organ transplant rejection, and in treating glomerulonephritis. Antibodies and analogs against L-selectin are useful in treating ischemia associated with reperfusion injury. Antibodies to PTHrp are useful in treating bone disease and metastatic cancer. In a particular embodiment, human antibodies against IL-8 may be used for the treatment or prevention of a pathology or condition

15

20

25

30

associated with IL-8. Such conditions include, but are not limited to, tumor metastasis, reperfusion injury, pulmonary edema, asthma, ischemic disease such as myocardial infarction, inflammatory bowel disease (such as Crohn's disease and ulcerative colitis), encephalitis, uveitis, 5 autoimmune diseases (such as rheumatoid arthritis, Sjögren's syndrome, vasculitis), osteoarthritis, gouty arthritis, nephritis, renal failure, dermatological conditions such as inflammatory dermatitis, psoriasis, vasculitic urticaria and allergic angiitis, retinal uveitis, conjunctivitis, 10 neurological disorders such as stroke, multiple sclerosis and meningitis, acute lung injury, adult respiratory distress syndrome (ARDS), septic shock, bacterial pneumonia, diseases involving leukocyte diapedesis, CNS inflammatory disorder, multiple organ failure, alcoholic hepatitis, antigen-antibody 15 complex mediated diseases, inflammation of the lung (such as pleurisy, aveolitis, vasculitis, pneumonia, chronic bronchitis, bronchiectasis, cystic fibrosis), Behcet disease, Wegener's granulomatosis, and vasculitic syndrome.

Typical autoimmune diseases which can be treated using the above-mentioned antibodies and analogs include systemic lupus erythematosus, rheumatoid arthritis, psoriasis, Sjogren's scleroderma, mixed connective tissue disease, dermatomyositis, polymyositis, Reiter's syndrome, Behcet's disease, Type 1 diabetes, Hashimoto's thyroiditis, Grave's disease, multiple sclerosis, myasthenia gravis and pemphigus.

For therapeutic applications, the antibodies may be administered in a pharmaceutically acceptable dosage form. They may be administered by any means that enables the active agent to reach the desired site of action, for example, intravenously as by bolus or by continuous infusion over a period of time, by intramuscular, subcutaneous, intraarticular, intrasynovial, intrathecal, oral, topical or inhalation routes. The antibodies may be administered as a single dose or a series of treatments.

For parenteral administration, the antibodies may be formulated as a solution, suspension, emulsion or

30

lyophilized powder in association with a pharmaceutically acceptable parenteral vehicle. If the antibody is suitable for oral administration, the formulation may contain suitable additives such as, for example, starch, cellulose, silica, various sugars, magnesium carbonate, or calcium phosphate. Suitable vehicles are described in the most recent edition of Remington's Pharmaceutical Sciences, A. Osol, a standard reference text in this field.

For prevention or treatment of disease, the appropriate dosage of antibody will depend upon known factors such as the pharmacodynamic characteristics of the particular antibody, its mode and route of administration, the age, weight, and health of the recipient, the type of condition to be treated and the severity and course of the condition, frequency of treatment, concurrent treatment and the physiological effect desired. The examples below are intended to illustrate but not to limit the invention.

In these examples, mice, designated XenoMouse™, are used for initial immunizations. A detailed description of the XenoMouse™ is found in the above referenced PCT application WO 94/02602. Immunization protocols appropriate to each antigen are described in the specific examples below. The sera of the immunized XenoMouse™ (or the supernatants from immortalized B cells) were titrated for antigen specific human antibodies in each case using a standard ELISA format. In this format, the antigen used for immunization was immobilized onto wells of microtiter plates. The plates were washed and blocked and the sera (or supernatants) were added as serial dilutions for 1-2 hours of incubation. After washing, bound antibody having human characteristics was detected by adding antihuman κ , μ , or γ chain antibody conjugated to horseradish peroxidase (HRP) for one hour. After again washing, the chromogenic reagent o-phenylene diamine (OPD) substrate and hydrogen peroxide were added and the plates were read 30 minutes later at 492 nm using a microplate reader.

Unless otherwise noted, the antigen was coated using plate coating buffer (0.1 M carbonate buffer, pH 9.6);

5

10

15

20

25

30

the assay blocking buffer used was 0.5% BSA, 0.1% Tween 20 and 0.01% thimerosal in PBS; the substrate buffer used in color development was citric acid 7.14 g/1; dibasic sodium phosphate 17.96 g/1; the developing solution (made immediately before use) was 10 ml substrate buffer; 10 mg OPD, plus 5 ml hydrogen peroxide; the stop solution (used to stop color development) was 2 M sulfuric acid. The wash solution was 0.05% Tween 20 in PBS.

10

15

20

25

30

5

Example 1

Human Antibodies Against Human IL-6

Three to five XenoMouse aged 8-20 weeks were agematched and immunized intraperitoneally with 50 µg human IL-6 emulsified in incomplete Freund's adjuvant for primary immunization and in complete Freund's adjuvant for subsequent injections. The mice received 6 injections 2-3 weeks apart. Serum titers were determined after the second dose and following each dose thereafter. Bleeds were performed from the retrobulbar plexus 6-7 days after injections. The blood was allowed to clot at room temperature for about 2 hours and then incubated at 4°C for at least 2 hours before separating and collecting the sera.

ELISAs were conducted as described above by applying 100 μ l/well of recombinant human IL-6 at 2 μ g/ml in coating buffer. Plates were then incubated at 4°C overnight or at 37°C for 2 hours and then washed three times in washing buffer. Addition of 100 μ l/well blocking buffer was followed by incubation at room temperature for 2 hours, and an additional 3 washes.

Then, 50 μ l/well of diluted serum samples (and positive and negative controls) were added to the plates. Plates were then incubated at room temperature for 2 hours and again washed 3 times.

After washing, 100 μ l/well of either mouse antihuman μ chain antibody conjugated to HRP at 1/2,000 or mouse antihuman κ chain antibody conjugated to HRP at 1/2,000, diluted in blocking buffer was added. After a 1

hour incubation at room temperature, the plates were washed 3 times and developed with OPD substrate for 10-25 minutes. 50 μ l/well of stop solution was then added and the results read on an ELISA plate reader at 492 nm. The dilution curves resulting from the titration of serum from XenoMouse^{γ} after 6 injections are shown in Figure 3. The data in Figure 3 show production of anti-IL-6 immunoreactive with antihuman κ and antihuman μ detectable at serum dilutions above 1:1,000.

10

15

20

25

5

Example 2

Human Antibodies Against Human TNFa

Immunization and serum preparation were conducted as described in Example 1 except that human recombinant TNF α (at 5 μ g per injection) was substituted for human IL-6.

ELISAs were conducted as described in Example 1 except that the initial coating of the ELISA plate employed 100 μ l/well recombinant human TNF α at 1 μ g/ml in coating buffer.

The dilution curves for serum from XenoMouse™ after 6 inductions obtained are shown in Figure 4. Again significant titers of human anti-TNFα binding were shown.

Serum titers for h γ , h μ , and h κ after one and two immunizations of the XenoMouse^M are shown in Table 1. When challenged with TNF- α , the XenoMouse^M switches isotypes from a predominant IgM response in the first immunization to an immune response with a large IgG component in the second immunization.

TABLE 2. Anti TNF-alpha serum titer responses of Xenomouse-2.

Bleed 1: after 2 immunizations Bleed 2: after 3 immunizations

ELISA 5 Serum titers Specific for TNF-alpha titer titer titer · (via hy) (via hu) (via har) XM₂ 10 1 bleed 1 500 3,000 1,500 bleed 2 10,000 8.000 15.000 2 bleed 1 200 3.000 500 bleed 2 2,700 5,000 1.000 3 bleed 1 <500 2.000 1,500 bleed 2 15,000 24,000 25,000 4 bleed 1 500 2.500 1.500 bleed 2 70,000 4,000 72,000 5 bleed 1 <500 2,500 1,500 bleed 2 1,000 10,000 7,000 6 bleed 1 1.000 13,000 4,500 bleed 2 10,000 24,000 25,000 7 bleed 1 <500 2,500 1,500 bleed 2 5,000 4,000 9,000 8 bleed 1 <500 1.000 500 bleed 2 2,700 5,000 9,000 9 bleed 1 200 6,000 4.000 bleed 2 40,000 80,000 80,000 10 bleed 1 200 2.000 500 bleed 2 15,000 8,000 60,000 20 11 bleed 1 1,500 1,000 1.500 bleed 2 24,000 2,700 72,000 12 bleed 1 200 2,000 1,000 bleed 2 10,000 4,000 25,000 13 bleed 1 500 30,000 500 bleed 2

15

2,000

4.000

12,000

Example 3

Human antibodies Against Human CD4

The human CD4 antigen was prepared as a surface protein using human CD4 on transfected recombinant cells as follows. Human CD4 consists of the extracellular domain of CD4, the transmembrane domain of CD4, and the cytoplasmic domain corresponding to residues 31-142, of the mature { chain of the CD3 complex. Human CD4 zeta (F15 LTR) as described in Roberts et al., Blood (1994) 84:2878 was introduced into the rat basophil leukemic cell line RBL-2H3, described F Callan, M., et al., Proc Natl Acad Sci USA (1993) 90:1 54 using the Kat high efficiency transduction described by Finer et al., Blood (1994) 83:43. Briefly, RBL-2H3 cells at 10 6 cells per well were cultured in 750 μ l DMEM-+ 20% FBS (Gibco) and 16 μ g/ml polybrene with an equal volume of proviral supernatant for 2 hours at 37°C, 5% CO. One ml of medium was removed and 750 μ l of infection medium and retroviral supernatant were added to each well and the cultures incubated overnight. The cells were washed and expanded in DMEM- + 10% FBS until sufficient cells were available for sorting. The CD4 zeta transduced RBL-2H3 cells were sorted using the FACSTAR plus (Becton Dickinson). cells were stained for human CD4 with a mouse antihuman CD4 PE antibody and the top 2-3% expressing cells were selected.

Immunizations were conducted as described in Example 1 using 1 X 10^7 cells per mouse except that the primary injection was subcutaneous at the base of the neck. The mice received 6 injections 2-3 weeks apart. Serum was prepared and analyzed by ELISA as described in Example 1 except that the initial coating of the ELISA plate utilized $100~\mu l$ per well of recombinant soluble CD4 at $2~\mu g/m l$ of coating buffer. The titration curve for serum from XenoMouse^M after 6 injections is shown in Figure 5. Titers of human anti-CD4 reactivity were shown at concentrations representing greater than those of 1:1,000 dilution.

- 22 -

5

10

15

20

25

30

Example 4

Human Antibodies Against Human L-selectin

The antigen was prepared as a surface displayed protein in C51 cells, a high expressing clone derived by transfecting the mouse pre-B cell 300.19 with LAM-1 cDNA (LAM-1 is the gene encoding L-selectin) (Tedder, et al., J. Immunol (1990) 144:532) or with similarly transfected CHO cells. The transfected cells were sorted using fluorescent activated cell sorting using anti-Leu-8 antibody as label.

The C51 and the transfected CHO cells were grown in DME 4.5 g/l glucose with 10% FCS and 1 mg/ml G418 in 100 mm dishes. Negative control cells, 3T3-P317 (transfected with gag/pol/env genes of Moloney virus) were grown in the same medium without G418.

Primary immunization was done by injection subcutaneously at the base of the neck; subsequent injections were intraperitoneal. 70-100 million C51 or transfected CHO cells were used per injection for a total of five injections 2-3 weeks apart.

Sera were collected as described in Example 1 and analyzed by ELISA in a protocol similar to that set forth in Example 1.

For the ELISA, the transfected cells were plated into 96 well plates and cell monolayers grown for 1-2 days depending on cell number and used for ELISA when confluent. The cells were fixed by first washing with cold 1 x PBS and then fixing solution (5% glacial acetic acid, 95% ethanol) was added. The plates were incubated at -25°C for 5 minutes and can be stored at this temperature if sealed with plate sealers.

The ELISA is begun by bringing the plates to room temperature, flicking to remove fixing solution and washing 5 times with DMEM medium containing 10% FCS at 200 μ l per well.

The wells were treated with various serum dilutions or with positive or negative controls. Positive control wells contained murine IgGl monoclonal antibody to human L-selectin.

5

15

25

30

The wells were incubated for 45 minutes and monolayer integrity was checked under a microscope. The wells were then incubated with antihuman κ chain antibody or antihuman μ chain antibody conjugates with HRP described in Example 1. The plates were then washed with 1% BSA/PBS and again with PBS and monolayer integrity was checked. The plates were developed, stopped, and read as described above. The results for serum from XenoMouse[™] are shown in Figures 6 and 7; human antibodies both to L-selectin and control 3T3 cells were obtained. However, the serum titers are higher for the L-selectin-expressing cells as compared to parental 3T3 cells. These results show the XenoMouse[™] produces antibodies specific for L-selectin with human μ heavy chain regions and human κ light chains.

The antisera obtained from the immunized XenoMouse[™] were also tested for staining of human neutrophils which express L-selectin. Human neutrophils were prepared as follows:

peripheral blood was collected from normal volunteers with 100 units/ml heparin. About 3.5 ml blood was layered over an equal volume of One-step Polymorph Gradient (Accurate Chemical, Westbury, NY) and spun for 30 minutes at 450 x g at 20°C. The neutrophil fraction was removed and washed twice in DPBS/2% FBS.

25

5

10

15

20

The neutrophils were then stained with either;

- (1) antiserum from XenoMouse™ immunized with C51
 cells (expressing L-selectin);
- (2) as a negative control, antiserum from a
 30 XenoMouse™ immunized with cells expressing human gp39.

The stained, washed neutrophils were analyzed by FACS. The results for antiserum from XenoMouse™ are shown in Figure 8.

35

These results show the presence of antibodies in immunized XenoMouse™ serum which contain fully human light chains immunoreactive with L-selectin. The negative control

antiserum from mice immunized with gp39 does not contain antibodies reactive against human neutrophils.

Example 5

5

10

15

20

25

30

35

Human Antibodies Against Human gp39

gp39 (the ligand for CD40) is expressed on activated human CD4 T cells. The sera of XenoMouse[™] immunized with recombinant gp39 according to this example contained fully human antibodies immunospecific for gp39.

The antigen consisted of stable transfectants of 300.19 cells or of CHO cells expressing gp39 cDNA cloned into the mammalian expression vector P1K1.HUgp39/IRES NEO as shown in Figure 9. CHO cells were split 1:10 prior to transfection in DMEM 4.5 g/l glucose, 10% FBS, 2 mM glutamine, MEM, NEAA supplemented with additional glycine, hypoxanthine and thymidine. The cells were cotransfected with the gp39 vector at 9 µg/10 cm plate (6 x 10⁵ cells) and the DHFR expressing vector pSV2DHFRs (Subranani et al., Mol Cell Biol (1981) 9:854) at 1 µg/10 cm plate using calcium phosphate transfection. 24 hours later the cells were split 1:10 into the original medium containing G418 at 0.6 mg/ml. Cells producing gp39 were sorted by FACS using an anti-gp39 antibody.

Mice grouped as described in Example 1 were immunized with 300.19 cells expressing gp39 using primary immunization subcutaneously at the base of the neck and with secondary intraperitoneal injections every 2-3 weeks. Sera were harvested as described in Example 1 for the ELISA assay. The ELISA procedure was conducted substantially as set forth in Example 1; the microtiter plates were coated with CHO cells expressing gp39 grown in a 100 mm dish in DMEM, 4.5 g/1 glucose, 10% FCS, 4mM glutamine, and nonessential amino acid (NEAA) solution for MEM (100X). On the day preceding the ELISA assay, the cells were trypsinized and plated into well filtration plates at 10⁵ cells/200 µl well and incubated at 37°C overnight. The positive controls were mouse antihuman gp39; negative controls were antisera from mice immunized with an antigen other than gp39. 50 µl of sample were used

for each assay. The remainder of the assay is as described in Example 1.

The dilution curves for the sera obtained after 4 injections from mice immunized with gp39 expressed on CHO cells are shown in Figure 10. As shown, the sera contained antihuman gp39 immunospecificity which is detectable with anti-human κ and anti-human μ chain antibodies coupled to HRP.

10 Example 6

5

15

20

25

30

35

Preparation of Human Mabs Against Tetanus Toxin
The antibodies prepared in this example were

secreted by hybridomas obtained by immortalizing B cells from xenomice immunized with tetanus toxin. The immunization protocol was similar to that set forth in Example 1 using 50 μ g tetanus toxin emulsified in complete Freund's adjuvant for intraperitoneal primary immunization followed by subsequent intraperitoneal injections with antigen incorporated into incomplete Freund's adjuvant. The mice received a total of 4 injections 2-3 weeks apart.

After acceptable serum titers of antitetanus toxin C (anti-TTC) were obtained, a final immunization dose of antigen in PBS was give 4 days before the animals were sacrificed and the spleens were harvested for fusion.

The spleen cells were fused with myeloma cells P3X63-Ag8.653 as described by Galfre, G. and Milstein, C. Methods in Enzymology (1981) 73:3-46.

After fusion the cells were resuspended in DMEM, 15% FCS, containing HAT supplemented with glutamine, pen/strep for culture at 37°C and 10% CO₂. The cells were plated in microtiter plates and maintained in HAT-supplemented medium for two weeks before transfer to HAT-supplemented medium. Supernatants from wells containing hybridomas were collected for a primary screen using an ELISA.

The ELISA was conducted as described in Example 1 wherein the antigen coating consisted of 100 μ l/well of tetanus toxin C (TTC) protein at 2 μ g/ml in coating buffer,

followed by incubation at 4°C overnight or at 37°C for two hours. In the primary ELISA, HRP-conjugated mouse antihuman IgM was used as described in Example 1. Two hybridomas that secreted anti-TTC according to the ELISA assay, clone D5.1 and clone K4.1 were used for further analysis.

As shown in Figure 11, clone D5.1 secretes fully human anti-TTC which is detectable using HRP-conjugated antihuman μ chain antibody and HRP-conjugated antihuman κ chain antibody. This is confirmed in Figure 11.

The antibody secreted by D5.1 did not immunoreact in ELISAs using TNFa, IL-6, or IL-8 as immobilized antigen under conditions where positive controls (sera from xenomice immunized with TNFa, IL-6 and IL-8 respectively) showed positive ELISA results.

The complete nucleotide sequence of the cDNAs encoding the heavy and light chains of the monoclonal were determined as shown in Figures 12 and 13. polyA mRNA was isolated from about 106 hybridoma cells and used to generate cDNA using random hexamers as primers. Portions of the product were amplified by PCR using the appropriate primers.

The cell line was known to provide human κ light chains; for PCR amplification of light chain encoding cDNA, the primers used were HKP1 (5'-CTCTGTGACACTCTCCTGGGAGTT-3') for priming from the constant region terminus and two oligos, used in equal amounts to prime from the variable segments; B3 (5'-GAAACGACACTCACGCAGTCTCCAGC-3').

For amplification of the heavy chain of the antibody derived form D5.1 (which contains the human μ constant region), MG-24VI was used to prime from the variable and μ P1 (5'-TTTTCTTTGTTGCCGTTGGGGTGC-3') was used to prime from the constant region terminus.

Referring to Figure 12 which sets forth the sequence for the heavy chain of the antibody secreted by clone D5.1, this shows the heavy chain is comprised of the human variable fragment VH6, the human diversity region DN1 and the human joining segment JH4 linked to the human μ constant region. There were two base-pair mutations from the germline sequence in the variable region, both in the CDRs.

5

10

15

20

25

30

Two additional mutations were in the D segment and six nongermline nucleotide additions were present at the D.-J. junction.

Finally, referring to Figure 13 which presents the light chain of the antibody secreted by D5.1, the human κ variable region B3 and human κ joining region JK3 are shown. There are nine base-pair differences from the germline sequences, three falling with CDR1.

10

15

20

25

5

Example 7

Human Antibodies Against PTHrp

Groups of XenoMouse^{N-2} were immunized intraperitoneally with either PTHrp (1-34) conjugated with BTG, as described by Ratcliffe et al., J. Immunol. Methods 127:109 (1990), or with PTHrp (1-34) synthesized as a 4 branched-MAP (multiple antigenic peptide system). The antigens were emulsified in CFA (complete Freunds adjuvant) and injected i.p. at a dose of 25 μ g per animal at 2 week intervals, and bled after two injections. The sera obtained from this bleed were analyzed by ELISA as described supra.

Serum titers for h γ , h μ , and h κ after one immunization of the XenoMouse[™] are shown in Table 2. When immunized with PTHrp, the XenoMouse[™] showed low serum titers in 5 of 7 mice on the first bleed, but when PTHrp-MAP is used, 7 of 7 mice show high serum titers on the first bleed.

TABLE 1. AntiPTHrp serum titer responses of Xenomouse-2.

First bleed after 2 immunizations with either PTHrp-BTG conjugate

10

15

20

25

30

35

<u>XM2</u>	Human Responses			
PTHrp-BTG Conjugate	titer (via hy)	titer (via h <i>µ</i>)	titer (via ha)	
1	<30	850	100	
2	<30	3.000	50	
3	<30	7,000	1,000	
4	<30	800	200	
5	<30	400	90	
6	<30	500	50	
· 7	<30	300	50	
XM2 PTHrp-MAP	titer (via h <i>y</i>)	titer (via h <i>µ</i>)	titer (via h <i>r</i>)	
1	<30	1,000	50	
2	<30	2,500	300	
3	<30	1,200	150	
4	150	1,000	270	
		2.500	300	
5	100	2,500	300	
5 6	<30	1,000	150	

Example 8

<u>Human Antibodies Against Human IL-8</u>

Immunization and serum preparation were as described in Example 1 except that human recombinant IL-8 was used as an immunogen.

ELISA assays were performed with respect to the recovered serum, also exactly as described in Example 1, except that the ELISA plates were initially coated using 100 μ l/well of recombinant human IL-8 at 0.5mg/ml in the coating buffer. The results obtained for various serum dilutions from XenoMouse[™] after 6 injections are shown in Figure 14.

Human anti-IL-8 binding was again shown at serum dilutions having concentrations higher than that represented by a 1:1,000 dilution.

5 Example 9

<u>Preparation of High Affinity Human Monoclonal Antibodies</u> Against Human IL-8

Groups of 4 to 6 XenoMouse™ aged between 8 to 10 weeks old were used for immunization and for hybridoma XenoMouse™ were immunized intraperitoneally with 25 μg of human recombinant-IL-8 (Biosource International, CA, USA) emulsified in complete Freund's adjuvant (CFA, Sigma) for the primary immunization. All subsequent injections were done with the antigen incorporated into incomplete Freund's adjuvant (IFA, Sigma). For animals used as spleen donors for hybridoma generation a final dose of antigen in phosphate buffer saline (PBS) was given 4 days before the fusion. Serum titers of immunized XenoMouse™ were first analyzed after a secondary dose of antigens, and from there after, following every antigen dose. Test bleeds were performed 6 to 7 days after the injections, by bleeding from the retrobulbar plexus. Blood was allowed to clot at room temperature for about 2 hours and then incubated at 4°C for at least 2 hours before separating and collecting the sera.

25

30

35

10

15

20

Generation of hybridomas

Spleen cells obtained from XenoMouse™ previously immunized with antigen, were fused with the non secretory NSO myeloma cells transfected with bcl-2 (NSO-bcl2) as described in Galfre G, et al., Methods in Enzymology 73, 3-46, (1981). Briefly, the fusion was performed by mixing washed spleen cells and myeloma cells at a ratio of 5:1 and gently pelleting them by centrifugation at 800Xg. After complete removal of the supernatant the cells were treated with 1 ml of 50% PEG/DMSO (polyethylene glycol MW 1500, 10% DMSO, Sigma) which was added over 1 min., the mixture was further incubated for one minute, and gradually diluted with 2 ml of DMEM over 2 minutes and diluted further with 8 ml of DMEM

over 3 minutes. The process was performed at 37 ºC with continued gentle stirring. After fusion the cells were resuspended in DMEM, 15% FCS, containing HAT, and supplemented with L glutamine, pen/strep, for culture at 37 °C and 10% CO2 in air. Cells were plated in flat bottomed 96 well microtiter trays. Cultures were maintained in HAT supplemented media for 2 weeks before transfer to HT supplemented media. Cultures were regularly examined for hybrid cell growth, and supernatants from those wells containing hybridomas were collected for a primary screen analysis for the presence of human μ , human gamma 2, and human kappa chains in an antigen specific ELISA as described above. Positive cultures were transferred to 48 well plates and when reaching confluence transferred to 24 well plates. Supernatants were tested in an antigen specific ELISA for the presence of human μ , human gamma 2, and human kappa chains.

As shown in Table 3 several hybridomas secreting fully human monoclonal antibodies with specificity for human IL-8 have been generated from representative fusions. In all of these human monoclonal antibodies the human gamma-2 heavy chain is associated with the human kappa light chain.

5

10

15

TABLE 3: ELISA determination of heavy and light chain composition of anti-IL-8 human monoclonal antibodies generated in XenoMouse^m

Sample ID	lg class	reactivity to hIL8			Total hlgG	
		titers	H _k OD (1:1)	mλ OD (1:1)	bγ OD (1:1)	(ng/ml)
Bkgd			0.08	0.04	0.12	
I8D1.1	hlgG2	500	4.12	0.04	4.09	1,159
18K2.1	hlgG2	200	4.18	0.18	4.11	2.000
18K2.2	hlgG2	1,000	4.00	0.04	4.00	4,583
I8K4.2	hlgG2	200	3.98	0.04	3.49	450
18K4.3	hlgG2	200	3.80	0.05	4.09	1,715
18K4.5	hlgG2	1,000	4.00	0.06	4.00	1,468

15

10

5

Evaluation of kinetic constants of XenoMouse™ hybridomas

In order to determine the kinetic parameters of these antibodies, specifically their on and off rates and their dissociation constants (KD), they were analyzed on the BIAcore instrument (Pharmacia). The BIAcore instrument uses plasmon resonance to measure the binding of an antibody to an antigen-coated gold chip.

25

30

20

BIAcore reagents and instrumentation:

The BIAcore instrument, CM5 sensor chips, surfactant P20, and the amine coupling kit containing N-hydroxysuccinimide (NHS), N-ethyl-N-(3-diethylaminopropyl)-carbodimide (EDC), and ethanolamine were purchased from

Immobilization of human recombinant Pharmaicia Biosensor. IL-8 onto the sensor surface was carried out at low levels of antigen density immobilized on the surface and was performed according to the general procedures outlined by the manufacturers. Briefly, after washing and equilibrating the instrument with HEPES buffer (HBS; 10 mM HEPES, 150 mM NaCl, 0.05% surfactant P20, pH 7.4) the surface was activated and IL-8 immobilized for the subsequent binding and kinetic The sensor surface was activated with 5 μ l of a studies. mixture of equal volumes of NHS (0.1 M) and EDC (0.1 M) injected at 10 μ l/min across the surface for activation, then 5 μ l of the ligand (human recombinant IL-8) at 12 μ g/ml in 5 mM maleate buffer, pH 6.0 was injected across the activated surface, and finally non-conjugated active sites were blocked with an injection of 35 μ l of 1 M ethanolamine. The surface was washed to remove non-covalently bound ligand by injection of 5 μ l 0.1 M HCI. All the immobilization procedure was carried out with a continuous flow of HBS of 10 μ l/min. About 100 resonance units (RU) of ligand (82 and 139 RU, in separate experiments) were immobilized on the sensorship, (according to the manufacturers 1,000 RU corresponds to about 1 ng/mm2 of immobilized protein).

These ligand coated surfaces were used to analyze hybridoma supernatants for their specific binding to ligand and for kinetic studies. The best regenerating condition for the analyte dissociation from the ligand in these sensorships was an injection of 10 μ l 100 mM HCl with no significant losses of binding observed after many cycles of binding and regeneration.

30

35

5

10

15

20

25

Determination of the dissociation, and association rates and the apparent affinity constants of fully human monoclonal antibodies specific for IL-8.

The determination of kinetic measurements using the BIAcore in which one of the reactants is immobilized on the sensor surface was done following procedures suggested by the manufacturers and described in Karlsson et al. "Kinetic analysis of monoclonal antibody-antigen interaction with a

new biosensor based analytical system." J. Immunol. Methods (19910 145, 229. Briefly the single site interaction between two molecules A and B is described by the following equation.

d[AB]/dt=ka[A][B]-kd[AB]

In which B is immobilized on the surface and A is injected at a constant concentration C. The response is a measure of the concentration of the complex [AB] and all concentration terms can be expressed as Response Units (RU) of the BIAcore:

dR/dt-kaC(Rmax-R) - kdR

where dR/dt is the rate of change of the signal, C is the concentration of the analyte, Rmax is the maximum analyte 15 binding capacity in RU and R is the signal in RU at time t. In this analysis the values of ka and kd are independent of the concentration of immobilized ligand on the surface of the The dissociation rates (kd) and association rates (ka) were determined using the software provided by the 20 manufacturers, BIA evaluation 2.1. The dissociation rate constant was measured during the dissociation phase that extended for 10 minutes at a constant buffer flow rate of 45 ul/min, after the completion of the injection of the 25 hybridoma supernatants onto the surface containing immobilized IL-8. The association phase extended over 1.25 minutes at a flow rate of 45 ul/min and the data was fitted into the model using the previously determined kd values. At least two surfaces with different levels of immobilized ligand were used in which different concentrations of anti 30 IL-8 hybridoma supernatants were tested for binding and analyzed for kinetic data. The kinetic constants determined on these two surfaces are presented in Table 4. affinities were determined to be very, ranging from 7 X 10.11 to 2 X 10.9 M. This compares vary favorably with the 35 affinities of murine monoclonal antibodies derived from normal mice.

5

TABLE 4: Kinetic constants of fully human monoclonal antibodies (lgG2, kappa) derived from XenoMouse™ II-a with specificity to human IL-8, determined by BIAcore.

Hybridoma	association rate ka (M ⁻¹ ,-1)	dissociation rate kd (-¹)	Dissociation Constant KD (M)=kd/ka	BIAcore surface h-IL-8 [RU]
I8D1-1	3.36 x 106	2.58 x 10-4	7.70 x 10-11	<u>81</u>
	2.80 x 106	1.73 x 10-4	6.20 x 10-11	134
I8K2-1	4.38 x 105	6.73 x 10-4	1.54 x 10-9	<u>81</u>
	3.83 x 105	6.85 x 10-4	1.79 x 10-9	134
18K2-2	5.24 x 105	2.26 x 10-4	4.30 x 10-10	<u>81</u>
	4.35 x 105	2.30 x 10-4	5.30 x 10-10	134
I8K4-2	5.76 x 106	8.17 x 10-4	1.42 x 10-10	<u>81</u>
	1.95 x 106	3.84 x 10-4	1.96 x 10-10	134
I8K4-3	2.66 x 106	7.53 x 10-4	2.83 x 10-10	<u>81</u>
	1.46 x 106	5.72 x 10-4	3.90 x 10-10	134
I8K4-5	4.00 x 105	9.04 x 10-4	2.26 x 10-9	<u>81</u>
	1.70 x 105	4.55 x 10-4	2.68 x 10-9	134

10

15

20

25

5

Methods for isolation of human neutrophils and assays for antibody activity

The primary in vivo function of IL-8 is to attract and activate neutrophils. Neutrophils express on their surface two distinct receptors for IL-8, designated the A receptor and the B receptor. In order to determine whether the fully human antibodies could neutralize the activity of IL-8, two different in vitro assays were performed with human neutrophils. In one assay, the ability of the antibodies to block binding or radiolabelled IL-8 to neutrophil IL-8 receptors was tested. In a second assay, the antibodies were tested for their ability to block an IL-8-induced

neutrophil response, namely the upregulation of the integrin Mac-1 on the neutrophil surface. Mac-1 is composed of two polypeptide chains, CD11b and CD18. Typically, anti-CD11b antibodies are used for its detection.

5

10

15

20

25

30

35

Isolation of neutrophils:

Human neutrophils are isolated from either freshly drawn blood or buffy coat. Human blood is collected by venipuncture into sterile tubes containing EDTA. Buffy coats are obtained from Stanford Blood Bank. They are prepared by centrifuging anticoagulated blood (up to 400 ml) in plastic bags at 2600 xg for 10 min at 20°C with the brake off. plasma supernatant is aspirated out of the bag and the buffy coat, i.e., the upper cell layer (40-50 ml/bag) is collected. One unit of buffy coat (40-50 ml) is diluted to final volume of 120 ml with Ca2, Mg2-free PBS. 30 milliliters of blood or diluted buffy coat are transferred into 50-ml centrifuge tubes on top of a 20-ml layer of Ficoll-Paque Plus (Pharmacia Biotech). The tubes are centrifuged at 500 xg for 20 min at 20°C with brake off. The supernatant, the mononuclear cells at the interface, and the layer above the pellet are carefully withdrawn. To completely remove the mononuclear cells, the cell pellet containing neutrophils and erythrocytes is resuspended with 5 ml of PBS and transferred into clean 50-ml tubes. The cells are washed in Ca2., Mg2.free PBS (300 xg for 5 min at 4°C). The erythrocytes are then lysed with ammonium chloride. The cells are resuspended in 40 ml of an ice-cold solution containing 155 mM NH,Cl and 10 nM EDTA, pH 7.2-7.4. The tubes are kept on ice for 10 min with occasional mixing and then centrifuged at 300 xg for 5 min at 4°C. The pellet is resuspended in PBS and washed once (300 xg for 5 min at 4°C). If erythrocyte lysis appears incomplete, the treatment with ammonium chloride is repeated. The neutrophils are again washed and finally suspended either in assay medium (RPMI-1640 supplemented with 10% fetal calf serum, 2 mM L-glutamine, 5x10⁻⁵ 2-mercapthoethanol, 1X nonessential amino acids, 1 mM sodium pyruvate and 10 mM Hepes) at a density of 3x107 cells/ml or in a binding buffer (PBS

containing 0.1% bovine serum albumin and 0.02% NaN_3), at a density of 6×10^6 cells/ml.

IL-8 receptor binding assay:

Multiscreen filter plates (96-well, Millipore, MADV 5 N6550) were pretreated with a PBS binding buffer containing 0.1% bovine serum albumin and 0.02% NaN, at 25°C for 2 hours. A final volume of 150 μ l, containing 4x10 5 neutrophils, 0.23 nM [125] -human-IL-8 (Amersham, IM-249) and varying concentrations of antibodies made up in PBS binding buffer, 10 was added to each well, and plates were incubated for 90 min at 4°C. Cells were washed 5 times with 200 μ l of ice-cold PBS, which was removed by aspiration. The filters were airdried, 3.5 ml of scintillation fluid was added (Beckman Ready Safe) and filters were counted on a Beckman LS6000IC counter. 15 The data obtained is presented as % specific bound [I125]-IL-8, which is calculated as the cpm in the presence of antibody divided by the cpm in the presence of PBS binding buffer only and multiplied by 100 (Figure 15). All six of the human anti-IL-8 monoclonals tested blocked IL-8 binding to human 20 neutrophils.

Neutrophil CD11b (Mac-1) expression assay:

Human IL-8 at a final concentration of 10 nM was preincubated with varying concentrations of monoclonal 25 antibodies at 4°C for 30 minutes and at 37°C for an additional 30 min. Neutrophils (4x105/well) were exposed to IL-8 in the presence or absence of antibodies at 4°C for 90 min, and incubated with PE-conjugated mouse-anti-human-CD11b (Becton Dickinson) for 45 min at 4°C. The cells were washed 30 with ice-cold PBS containing 2% fetal calf serum. Fluorescence was measured on a Becton Dickinson FACscan cell analyzer. A mouse monoclonal antibody against human CD11b obtained from R&D System, Inc. was used as a positive control while the purified myeloma human IgG2 (Calbiochem) was used 35 as a negative control in the experiments. The expression levels of CD11b on neutrophils were measured and expressed as the mean fluorescence channel. The mean fluorescence channel

derived form the negative control antibody was subtracted from those of experimental samples.

5	% inhibition =	only	mean fluorescence the presence of antibodies	
10	* Innibition =	mean fluorescence in the presence of IL-8 only	mean fluorescence	

As shown in Table 5, five of the six antibodies blocked upregulation of CD11b to some degree, with three of the five giving complete blocking.

TABLE 5: Inhibition of CD11b expression on human neutrophils by monoclonal antibodies against IL-8.

20	Antibody	Concentration (nM)	Inhibition of CD11b expression (%)
	R&D anti-IL8	333	100
	I8K1.1	6	100
	I8K2.1	10	60
	I8K2.2	. 32	100
25	I8K4.2	3	10
	I8K4.3	8	100
	I8K4.5	5	0
	Human IgG2	33	0
	·		

Background of CD11b expression is 670 (mean fluorescence) while CD11b expression in the presence of 10 nM of human IL-8 is 771.

<u>Sequence analysis of Immunoqlobulin transcripts derived from anti-hIL-8 hybridomas.</u>

All sequences were derived by direct sequencing of PCR fragments generated form RT-PCR reactions of RNA prepared from hybridomas D1.1, K2.2, K4.2 and K4.3, using human V_z and human V_z family specific primers (Marks et. al. 1991; Euro J. Immunol 21;985-991) and a primer specific for either the human gamma 2 constant region (MG-40d; 5'GCTGAGGGAGTAGAGTCCTGAGGACTGT-3') or human kappa constant

region (HKP2; Green et al 1994; Nature Genetics 7: 13-21)).

In Figure 16 A-H, both strands of the four clones were sequenced and analyzed to generate the complete sequence.

All sequences were analyzed by alignments to the "V BASE sequence directory", Tomlinson et al., MRC Centre for Protein Engineering, Cambridge, UK. The variable and joining regions are indicated by brackets []. Nucleotides containing an "N" indicate uncertainty in the generated sequence.

Based on sequence alignments with sequences found in the V-base database the heavy chain transcript from hybridoma D1.1 has a human $V_H4-21\,(DP-63)$ variable region (7 point mutations were observed compared to the germline sequence), a human 21-10rc D segment, a human J_H3 joining region and a human gamma 2 constant region. See Figure 16A.

The kappa light chain transcript from hybridoma

25 D1.1 is comprised of a human kappa variable region with homology to V, 08/018 (DPK1) (16 point mutations were observed when compared to the germline sequence) a human J,3 joining region, and a human kappa constant region. See Figure 16B.

Based on sequence alignments with sequences found in the V-base database the heavy chain transcript from hybridoma K2.2 has a human $V_{\rm H}3-30$ variable region (3 point mutations were observed compared to the germline sequence), a human IR3rc D segment, a human $J_{\rm H}4$ joining region and a human gamma 2 constant region. See Figure 16C.

The kappa light chain transcript from hybridoma K2.2 is comprised of a human kappa variable region with homology to V_kIV (B3; DPK24) (9 point mutations were observed

5

10

15

when compared to the germline sequence), a human J,3 joining region, and a human kappa constant region. See Figure 16D.

Based on sequence alignments with sequences found in the V-base database the heavy chain transcript from hybridoma K4.2 has a human $V_{\rm H}4-34$ variable region (8 point mutations were observed compared to the germline sequence), a human K1 D segment, a human $J_{\rm H}4$ joining region and a human gamma 2 constant region. See Figure 16E.

The kappa light chain transcript from hybridoma K4.2 is comprised of a human kappa variable region with homology to V, 08/018 (DPK1) (6 point mutations were observed when compared to the germline sequence), a human J,4 joining region, and a human kappa constant region. See Figure 16F.

Based on sequence alignments with sequences found in the V-base database the heavy chain transcript from hybridoma K4.3 has a human $V_{\rm H}5-51$ (DP-73) variable region, a human M5-a/M5-b D segment, a human $J_{\rm H}4$ joining region and a human gamma 2 constant region. See Figure 16G.

The kappa light chain transcript from hybridoma K4.3 is comprised of a human kappa variable region with homology to V, 02/012 (DPK9) (9 point mutations were observed when compared to the germline sequence), a human J,4 joining region, and a human kappa constant region. See Figure 16H.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

5

10

15

20

25

30

Biological Deposits

5

10

15

yH1C contained in *S. cerivisiae* was deposited with the American Type Culture Collection ("ATCC"), 12301 Parklawn Drive, Rockville MD 20852, USA, on April 26, 1996, and given ATCC accession no. ______. The deposit of this YAC is for exemplary purposes only, and should not be taken as an admission by the Applicant that such deposit is necessary for enablement of the claimed subject matter.

In respect of all designated States in which such action is possible and to the extent that it is legally permissible under the law of the designated State, it is requested that a sample of the deposited micro-organism be made available only by the issue thereof to an independent expert, in accordance with the relevant patent legislation, e.g., EPC rule 28(4), United Kingdom Patent Rules 1982 rule 17(3), Australian Regulation 3.25(3) and generally similar provisions mutatis mutandis for any other designated State.

International Application No: PCT'

MICROORGANISMS
Optional Sheet in connection with the microorganism referred to on page 41 , lines $1-20$ of the description
A. IDENTIFICATION OF DEPOSIT
Further deposits are identified on an additional sheet '
Name of depositary insulation.
American Type Culture Collection
Address of depositary institution (including postal code and country)
12301 Parklawn Drive Rockville, MD 20852 US
Date of deposit * April 26, 1996 Accession Number
B. ADDITIONAL INDICATIONS ' Heave blank if not applicable). This intermation is continued on a separate anached short
C. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE
D. SEPARATE FURNISHING OF INDICATIONS (Ideave blank if not applicable)
The indications listed below will be submitted to the International Burasu later [†] (Specify the general nature of the indications e.g., "Accession Number of Debosit")
E. W This sheet was received with the International application when filed (to be checked by the receiving Office) (Authorized Officer)
The date of receipt (from the applicant) by the International Bureau
was (Authorized Officer)

Form PCT/RO/134 (January 1981)

CLAIMS

1. A method to produce a human immunoglobulin or an analog thereof, specific for a desired antigen, which method comprises:

administering said antigen or an immunogenic portion thereof to a nonhuman animal under conditions to stimulate an immune response, whereby said animal produces B cells that secrete immunoglobulin specific for said antigen; wherein said nonhuman animal is characterized by being substantially incapable of producing endogenous heavy and light immunoglobulin chains, but capable of producing human immunoglobulin; and

recovering said immunoglobulin or analog.

15

10

5

- 2. The method of claim 1 wherein said recovering step comprises recovering polyclonal immunoglobulin or analog from said animal.
- 3. The method of claim 1 wherein said recovering step comprises immortalizing B cells from said animal immunized with said antigen, screening the resulting immortalized cells for the secretion of said immunoglobulin specific for said antigen, and
- a) recovering immunoglobulin secreted by said immortalized B cells, or
 - b) recovering the genes encoding at least the immunoglobulin from the immortalized B cells, and optionally modifying said genes;
- expressing said genes or modified forms thereof to produce immunoglobulin or analog; and recovering said immunoglobulin or analog.
- 4. The method of claim 1 wherein said recovering 35 step comprises:

recovering genes encoding the immunoglobulins from the primary B cells of the animal;

generating a library of said genes expressing the immunoglobulins;

screening the library for an immunoglobulin with the desired affinity for the antigen;

recovering the genes encoding the immunoglobulin; expressing said recovered genes to produce an immunoglobulin or analog recovering said immunoglobulin or analog.

- 5. A recombinant DNA molecule comprising a nucleotide sequence encoding the immunoglobulin or analog produced by the method of claim 1.
- 6. The DNA molecule of claim 5 wherein said encoding-nucleotide sequence is operably linked to control sequences capable of effecting its expression.
 - 7. A cell or cell line modified to contain the DNA molecule of claim 6.

20

25

5

- 8. A method to produce a fully human immunoglobulin or an analog thereof which method comprises culturing the cells of claim 7 under conditions whereby said encoding nucleotide sequence is expressed to produce said immunoglobulin or analog; and recovering said immunoglobulin or analog.
- 9. A DNA molecule comprising a nucleotide sequence corresponding to the gene or modified gene prepared by the method of claim 3.
 - 10. The DNA molecule of claim 9 wherein said encoding nucleotide sequence is operably linked to control sequences capable of effecting its expression.

35

11. A cell or cell line modified to contain the DNA molecule of claim 9.

12. A method to produce a fully human immunoglobulin or an analog thereof which method comprises culturing the cells of claim 11 under conditions whereby said encoding nucleotide sequence is expressed to produce said immunoglobulin or analog; and recovering said immunoglobulin or analog.

- 13. A DNA molecule which comprises a nucleotide sequence encoding a human immunoglobulin with desired
 10 affinity prepared according to the method of claim 4.
 - 14. The DNA molecule of claim 13 wherein said encoding nucleotide sequence is operably linked to control sequences capable of effecting its expression.
 - 15. A cell or cell line modified to contain the DNA molecule of claim 13.
- immunoglobulin or an analog thereof which method comprises culturing the cells of claim 15 under conditions whereby said encoding nucleotide sequence is expressed to produce said immunoglobulin or analog; and recovering said immunoglobulin or analog.
 - 17. An immortalized B cell which secretes a fully human immunoglobulin to a desired antigen prepared according to claim 3.
- 18. A method to produce an immunoglobulin or analog which comprises culturing the cells of claim 17 and recovering said immunoglobulin or analog.
- 19. A fully numan immunoglobulin or analog35 produced by the method of claim 1.

5

15

20. The immunoglobulin or analog of claim 19 wherein the desired antigen is selected from the group consisting of

the leukocyte markers, CD2, CD3, CD4, CD5, CD6, CD7, CD8, CD11a,b,c, CD13, CD14, CD18, CD19, CD20, CD22, CD23, CD27 and its ligand, CD28 and its ligands B7.1, B7.2, B7.3, CD29 and its ligand, CD30 and its ligand, CD40 and its ligand gp39, CD44, CD45 and isoforms, CDw52 (Campath antigen), CD56, CD58, CD69, CD72, CTLA-4, LFA-1 and TCR;

the histocompatibility antigens, MHC class I or II, the Lewis Y antigens, SLex, SLey, SLea, and SLeb;

the integrins, VLA-1, VLA-2, VLA-3, VLA-4, VLA-5, VLA-6, α V β 3, and LFA-1, Mac-1, and pl50,95, $\alpha_{\nu}\beta_{1}$, gpIIbIIIa, $\alpha_{R}\beta_{1}$, $\alpha_{6}\beta_{4}$, $\alpha_{\nu}\beta_{5}$, $\alpha_{\nu}\beta_{6}$, and $\alpha_{\nu}\beta_{7}$;

the selectins, L-selectin, P-selectin, and E-selectin and their counterreceptors VCAM-1, ICAM-1, ICAM-2, and LFA-3;

the interleukins, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, and IL-15;

the interleukin receptor is selected from the group consisting of IL-1R, IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R, IL-8R, IL-9R, IL-10R, IL-11R,, IL-12R, Il-13R,, IL-14R, and IL-15R;

25 the chemokine is selected from the group consisting of PF4, RANTES, MIP1α, MCP1, NAP-2, Groα, Groβ, and IL-8;

the growth factor is selected from the group consisting of TNFalpha, TGFbeta, TSH, VEGF/VPF, PTHrP, EGF family, FGF, PDGF family, endothelin, Fibrosin (F_sF_{-1}) , human Laminin, and gastrin releasing peptide (GRP);

the growth factor receptor is selected from the group consisting of TNFalphaR, RGFbetaR, TSHR, VEGFR/VPFR, FGFR, EGFR, PTHrPR, PDGFR family , EPO-R, GCSF-R and other hematopoietic receptors;

the interferon receptor is selected from the group consisting of IFNC α R, IFN β R, and IFN λ R;

the Ig and its receptor is selected from the group consisting of IgE, FceRI, and FCERII;

20

the tumor antigen is selected from the group consisting of her2-neu, mucin, CEA and endosialin;

the allergen is selected from the group consisting of house dust mite antigen, lol pl (grass) antigens, and urushiol;

the viral protein is selected from the group consisting of CMV glycoproteins B, H, and gCIII, HIV-1 envelope glycoproteins, RSV envelope glycoproteins, HSV envelope glycoproteins, HPV envelope glycoproteins, Hepatitis family surface antigens;

the toxin is selected from the group consisting of pseudomonas endotoxin and osteopontin/uropontin, snake venom, spider venom, and bee venom conotoxin;

the blood factor is selected form the group

consisting of complement C3b, complement C4a, complement C4b
Rh factor, fibrinogen, fibrin, and myelin associated growth inhibitor; and

the enzyme is selected from the group consisting of cholesterol ester transfer protein, membrane bound matrix metalloproteases, and glutamic acid decarboxylase (GAD).

- 21. The immunoglobulin or analog of claim 14 wherein said desired antigen is selected from the group consisting of human IL-6, human IL-8, human TNF α , human CD4, human L-selectin, human gp39, human IgE, human $\alpha V\beta$ 3, human Fibrosin (F,F.₁), human laminin, human PTHrp, and tetanus toxin C(TTC).
- 22. A recombinant DNA molecule comprising a nucleotide sequence that encodes the immunoglobulin or analog of claim 19-21.
- 23. The DNA molecule of claim 22 wherein said encoding nucleotide sequence is preferably linked to control sequences capable of effecting its expression.
 - 24. A cell or cell line modified to contain the DNA molecule of claim 23.

5

10

20

25. A method to produce an immunoglobulin or analog specific for a desired antigen which method comprises culturing the cell or cell line of claim 24 under conditions wherein said nucleotide sequence is expressed to produce said immunoglobulin or analog; and recovering the immunoglobulin or analog.

- 26. An human antibody or analog thereof which is specifically immunoreactive with an antigen selected from the group consisting of transition state mimics; leukocyte markers; histocompatibility antigens; adhesion molecules; interleukins; interleukin receptors; chemokines; growth factors; growth factor receptors; interferon receptors; Igs and their receptors, tumor antigens; allergens; viral proteins; toxins; blood factors; enzymes; and the miscellaneous antigens ganglioside GD3, ganglioside GB2, LMP1, LMP2, eosinophil major basic protein, eosinophil cationic protein, pANCA, Amadori protein, Type IV collagen, glycated lipids, λ-interferon, A7, P-glycoprotein, Fas (AFO-1) and oxidized-LDL.
- The antibody or analog of claim 26 wherein the leukocyte marker is selected from the group consisting of CD2, CD3, CD4, CD5, CD6, CD7, CD8. CD11a,b,c, CD13, CD14, CD18, CD19, CD20, CD22, CD23, CD27 and its ligand, CD28 and its ligands B7.1, B7.2, B7.3, CD29 and its ligand, CD30 and its ligand, CD40 and its ligand gp39, CD44, CD45 and isoforms, CDw52 (Campath antigen), CD56, CD58, CD69, CD72, CTLA-4, LFA-1 and TCR;
- the histocompatibility antigen is selected from the group consisting of MHC class I or II, the Lewis y antigens, SLex, SLey, Slea, and SLeb;

the adhesion molecule is selected from the group consisting of VLA-1, VLA-2, VLA-3, VLA-4, VLA-5, VLA-6, α V β 3, and LFA-1, Mac-1, pl50,95, $\alpha_{\nu}\beta_{1}$, gpIIbIIIa, $\alpha_{R}\beta_{3}$, $\alpha_{6}\beta_{4}$, $\alpha_{\nu}\beta_{5}$, $\alpha_{\nu}\beta_{6}$, and $\alpha_{\nu}\beta_{7}$, L-selectin, P-selectin, and E-selectin and their counterreceptors VCAM-1, ICAM-1, ICAM-2, and LFA-3;

5

10

15

20

25

30

the interleukin is selected from the group consisting of IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, ad IL-15;

the interleukin receptor is selected from the group consisting of IL-1R, IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R,

IL-8R, IL-9R, IL-10R, IL-11R, IL-12R, IL-13R, IL-14R, and IL-15R,

the chemokine is selected from the group consisting of PF4, RANTES, MIP1α, MCP1, NAP-2, Groα, Groβ, and IL-8; the growth factor is selected from the group consisting of TNFalpha, TGFbeta, TSH, VEGF/VPF, PthrP, EGF family, FGF, PDGF family, endothelia, Fibrosin (F,F.1), human Laminin, and gastrin releasing peptide (GRP);

the growth factor receptor is selected from the group consisting of TNFalphaR, RGFbetaR, TSHR, VEGFR/VPFR, FGFR, EGFR, PTHrPR, PDGFR family, EPO-R, GCSF-R and other hematopoietic receptors;

the interferon receptor is selected from the group consisting of IFN α R, IFN β R, and IFN γ R;

the Ig and its receptor is selected from the group IgE, FceRI, and FCeRII;

tumor antigen is selected from the group her2-neu, mucin, CEA and endosialin;

the allergen is selected from the group consisting of house dust mite antigen, lol pl (grass) antigens, and urushiol;

the viral protein is selected from the group consisting of CVM glycoproteins B, H, and GCIII, HIV-1 envelope glycoproteins, RSV envelope glycoproteins, HSV envelope glycoproteins, VZV envelope glycoproteins, HPV envelope glycoproteins, Hepatitis family surface antigens;

the toxin is selected form the group consisting of pseudomonas endotoxin and osteopontin/uropontin, snake venom, and bee venom;

the blood factor is selected from the group consisting of complement C3b, complement C5a, complement C5b-

5

20

9, RH factor, fibrinogen, fibrin, and myelin associated growth inhibitor; and

The enzyme is selected from the group consisting of cholesterol ester transfer protein, membrane bound matrix metalloproteases, and glutamic acid decarboxylase (GAD)

- 28. The antibody or analog of claim 26 wherein the desired antigen is selected from the group consisting of human IL-6, human IL-8, human TNFα, human CD4, human L-selectin, human gp39, human IgE and tetanus toxin C(TTC).
- 29. The antibody or analog of claim 19 wherein the desired antigen is human IL-6.
- 30. The antibody or analog of claim 19 wherein the described antigen is human IL-8.
 - 31. The antibody or analog of claim 19 wherein the desired antigen is human $TNF\alpha$.
 - 32. The antibody or analog of claim 19 wherein the desired antigen is human CD4.
- 33. The antibody or analog of claim 19 wherein the desired antigen is human L-selectin.
 - 34. The antibody or analog of claim 19 wherein the desired antigen is human gp39.
- 35. The antibody or analog of claim 19 wherein the desired antigen is tetanus toxin C (TTC).
 - 36. The antibody or analog of claim 19 wherein the desired antigen is human IgE.
 - 37. The antibody or analog of claim 19 wherein the desired antigen is human $\alpha V \beta 3$.


35


5

10

38. The antibody or analog of claim 19 wherein the desired antigen is human fibrosin.

- 39. The antibody or analog of claim 19 wherein the desired antigen is human PTHrp.
 - 40. The antibody or analog of claim 26 which is an agonist or is a catalyst.
- 10 41. A recombinant DNA molecule encoding the antibody of any of claim 26-40.
- 42. A recombinant DNA molecule which comprises an expression system for the antibody or analog of any claims
 26-40 which expression system comprises a nucleotide sequence encoding said antibody or analog operably linked to control sequences capable of effecting its expression.
- 43. A recombinant host cell which is modified to contain the DNA molecule of claim 42.
 - 44. A method to produce an antibody or analog which method comprises culturing cells of claim 43 under conditions wherein said coding sequence is expressed; and recovery the antibody of analog produced.








FIG.3




FIG.4








FIG.5









F 16.9




FIG. 10




FIG. 11
SUBSTITUTE SHEET (RULE 26)

		<u>CDR1</u>	_				
Germline JH	5. 1.4	AGACCCTCTC AGACCCTCTC	ACTCACCTGT ACTCACCTGT	GCCATCTCCG	GGGACAGTG	CTCTAGCAA	C 50 C 50
Germline D(Germline hW	'						-
Germine in	n u	-		20		VH6 ——	-
			CD	R2			
Germline VH Hybridomo D5 Germline JH Germline D(1,4 14	AGTGCTGCTT AGTGCTGCTT	GGAACTGGAT GGAACTGGAT	CAGGCAGTCC CAGGCAGTCC	CCATCGAGAG CCATCGAGAG	GCCTTGAGTG	100 100
Germline by	N1)					·	-
Germinine ima	ıu		v	H6 —			•
	•		▼.	110			-
Germline VH Hybridomo D5 Germline JH Germline D(1	. 1.4	GCTGGGAAGG GCTGGGAAGG	ACATACTACA ACATACTACA	GGTCCAAGTG GGTCCAAGTG	GTATAATGAT GTATAATGAT	TATGCAGTAT TATGCAGTUT	150 150
Germline hM	•						
			Vi	16 ——			
Germline VH6 Hybridoma D5 Germline JH4	. 1.4	CTGTGAAAAG CTGTGAAAAG	TCGAATAACC TCGAATAACC	ATCAACCCAG ATCAACCCAG	ACACATCCAA ACACATCCAA	GAACCAGTTC GAACCAGTTC	200 200
Germline D(N							
Germline hMu							
			VI	16 ———			
Germline VH6 Hybridoma D5. Germline JH4	1.4	TCCCTGCAGC TCCCTGCAGC	TGAACTCTGT TGAACTCTGT	GACTCCCGAG GACTCCCGAG	GACACGGCTG GACACGGCTG	TGTATTACTG TGTATTACTG	250 250
Germline D(N	•	· · · · · · · · · · · · · · · · · · ·					
Germline hMu							
	•		——— VH	b ———		-	
Germline VH6 Hybridomo D5. Germline JH4	1.4	IGCAAGAGA IGCAAGAGAT /	ATAGCAGITOC	CTGGCGTCCT		TGGGGCCAGG	
Germline D(N		T	TAGCAGCAG	CTGG	CTTGACTAGC	IGGGGCCAAG	20 15
Germline hMu	•		 -				13
	-	— VH6 ——	DN	1- -	JH4 ·		

FIG.12A

Germline Hybridomo Germline Germline	D5. 1.4 JH4		T CACCGTCTCC	C TCAGGGAGTO	CATCCGCCC	AACCCTTTT	C 350 - 43
Germline		JI		GCGAGT(CATCCCCCC	AACCCTTTT	27
Germline	D5. 1.4 JH4	CCCCTCGTCT	CCTGTGAGAA	TTCCCCGTCG	GATACGAGCA		259 400 43
Germline Germline	D(N1) hMu	CCCCTCGTCT		TTCCCCGTCG	GATACGAGCA	GCGTGGCCGT	15 77

FIG.12B

Germline B3 Hybridomo D5 1.4 Germline JK3 Germline CK		TGACCCAGTO	TCCAGACTC	CTGGCTGTG	CTCTGGGCGA
Germline B3 Hybridoma D5 1.4 Germline JK3 Germline CK	GAGGGCCACC ——ACC	ATCAACTGCA ATCAAGTGCA	AGTCCAGCCA AGTCCAGCCA	GAGTGTTTT	
	-		—— B3	3——	
Germline B3 Hybridoma D5 1.4 Germline JK3 Germline CK	PCAATAAGAA BCAATAAGAA	CTACTTAGCT CTACTTAGCT	TGGTACCAGO TGGTACCAGO	AGAAACCAGG AGAAACCAGG	ACAGCCTCCT ACAGCCTCCT
Dermittle Ch			— B3 —		
Germline B3 Hybridoma D5 1.4 Germline JK3	AACCTOCTCA AAACTACTCA	TTTACTGGGC TTTACTGGGC	ATCTACCCGG ATCTACCCGG	GAATCCGGGG GAATCCGGGG	TCCCTGACCG TCCCTGACCG
Germline CK			B3		
Germline B3 Hybridoma D5 1.4 Germline JK3 Germline CK		AGCGGGTCTG AGCGGGTCTG	GGACAGATTT GGACAGATTT	CACTCTCACC CACTCTCACC	ATGAGCAGCC ATGGCAGCC
oermi ille CA			B3		
Germline B3 Hybridoma D5 1.4 Germline JK3	TGCAGGCTGA TGCAGGCTGA	AGATGTGGCA AGATGTGGCA	GTTTATTACT GTTTATTACT	GTCAGCAATA GTCAGCAATA	TTATACTACT TTATACTATT
Germline CK			B3		
Hybridomo D5 1.4	CC————————————————————————————————————				
Germline CK		JK			GAACTGTGGC CK

FIG. 13A

. 4	TGCACCATCT	GTCTTCATCT	TCCCGCCATC	TGATGAGCAG	TTGAAATCT
					
			- CIV —		
. 4	GAACTGCCTC	TGTTGTGTGC	CTGCTGAATA	ACTICTATCC	CAGAGAGGCC
	GAACTGCCTC	TGTTGTGTGC	CTGCTGAATA	ACTICIATCO	CAGAGAGGCC
			—— СК —		
					
	AAAGTACAGT				
	. 4	TGCACCATCT	TGCACCATCT GTCTTCATCT 4 GAACTGCCTC TGTTGTGTGC GAACTGCCTC TGTTGTGTGC 4 AAAGTACAGT GGAAGGTGGA AAAGTACAGT GGAAGGTGGA	TGCACCATCT GTCTTCATCT TCCCGCCATC CK 4 GAACTGCCTC TGTTGTGTGC CTGCTGAATA GAACTGCCTC TGTTGTGTGC CTGCTGAATA CK 4 AAAGTACAGT GGAAGGTGGA TAACGCCCTC AAAGTACAGT GGAAGGTGGA TAACGCCCTC	TGCACCATCT GTCTTCATCT TCCCGCCATC TGATGAGCAG TGCACCATCT GTCTTCATCT TCCCGCCATC TGATGAGCAG CK GAACTGCCTC TGTTGTGTGC CTGCTGAATA ACTTCTATCC GAACTGCCTC TGTTGTGTGC CTGCTGAATA ACTTCTATCC CK 4 AAAGTACAGT GGAAGGTGGA TAACGCCCTC CAATCGGGTT AAAGTACAGT GGAAGGTGGA TAACGCCCTC CAATCGGGT- CK

FIG.13B






FIG. 14





15/18

FIG. 16A

[CAGTCTCCATCCTCCCTGTCTGCATCTGTAGGCGACAGAGTCACCATCACTTGCCAGGCGAGTC
AGGACATTAGTAAGTTTTTAAGTTGGTTTCAACAGAAACCAGGGAAAGCCCCTAAACTCCTGATC
TACGGTACATCCTATTTGGAAACCGGGGTCCCATCAAGTTTCAGTGGAAGTGGATCTGGGACAGA
TTTTACTCTCACCATCAGCAGCCTGCAGCCTGAAGATGTTGCAACATATTTCTGTAACAGNATG
ATGATCTCCC][ATACACTTTCGGCCCTGGGACCAAAGTGGATATCAAAC]GAACTGTGGCTGCAC
CATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAACTGCCTCTGTTGTGTGCC
TGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAACGCCC

FIG. 16B

16/18

[AGGTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTCACCTTCAGTAGCTATGGCATGCACTGGNT CCGCCAGGCTCCAGGCAAGGGGCTGGAGTGGGTGGCAGAAATATCATATGATGGAAGTAATAAA TACTATGTAGACTCCGTGAAGGGCCGACTCACCATCTCCAGAGACAATTCCAAGAACACGCTGT ATCTGCAAATGAACAGCCTGAGAGCTGAGGACACGGCTGTGTATTACTGTGCGAGAGA][CCGAC TGGGGAT][CTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCAG]CCTCCACCAAGG GCCCATCGGTCTCCCCCCTGGCGCCCTGCTCCAGGAGCACCTCCGAGAGCACAGC(GC)GGCCCT GGGCTGCCTGGTCCCCCGAGAGCACCGGTGTCGTGGAACTCAGGCGCTC TGACCAG

FIG. 16C

[CTGACNCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGAGGGCCACCATCAACTGCAAGT
CCAGCCAGAGTGTTTTATACATCTCCAACAATAAAACTACTTAGCTTGGTACCAGCAGAAACCA
GGACAGTCTCCTAAACTGCTCATTTACTGGGCATCTACCCGGAAATCCGGGGTCCCTGACCGATT
CAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGGCTGAAGATGTG
GCAGTTTATTACTGTCAACAGTATTATGATACTCC][ATTCACTTTCGGCCCTGGGACCAAAGTGG
ATATCAAAC]GAACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAA
TCTGGAACTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTG
GAAGGTGGNTAACGCCCCA

FIG. 16D

FIG. 16E

[TGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCAGGC
GAGTCAGGACATTAGTAACTATTTAAATTGGTATCAACAGAAAGCAGGGAAAGCCCCTAAGGTCC
TGATCTACGCTGCATCCAATTTGGAAGCAGGGGTCCCATCAAGGTTCAGTGGAAGTTGGATCTGGG
ACAGATTTTACTTTCACCATCAGCAGCCTGCAGCCTGAAGATATTGCAACATATTATTGTCAACA
CTATGATAATCT]A[CTCACTTTCGGCGGAGGGACCAAGGTAGAGATCAAAC]GAACTGTGGCTGC
ACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGACTGCCTCTGTTGTGTG
CCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGG

FIG. 16F

18/18

AGTCTCTGAAGATCTCCTGTAAGGGTTCTGGATACAGCTTTACCAGCTACTGGATCGGCTGGGTG
CGCCAGATGCCCGGGAAAGGCCTGGAGTGGATGGGGATCATCTATCCTGGTGACTCTGATACCA
GATACAGCCCGTCCTTCCAAGGCCAGGTCACCATCTCAGCCGACAAGTCCATCAGCACCGCCTA
CCTGCAGTGGAGCAGCCTGAAGGCCTCGGACACCGCCATGTATTACTGTGCGAGACA][GGACGG
TG][ACTCCTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCAG]CCTCCACCAAGGG
CCCATCGGTCTTCCCCCTGGCGCCCCTGCTCCAGGAGCACCTCCGAGAGCACAGC(GC)GGCCCTG
GGCTGCCTGGTCCAAGGACTACTTCCCCCGAACCGGTGACGGTGTCGTGGAACTCAGGCGCTCT
GACCAGCGGCGTGCACACCTTCCCACTGCCA

FIG. 16G

TGTCTGCATCTATTGGAGACAGAGTCACCATCACTTGCCGGGCAAGTCAGAGCATTAGCAACTA
TTTAAATTGGTATCAGCAGAAACCAGGGCAAAGCCCCTAAGTTCCTGATCTATGGTGCATCCAGT
TTGGAAAGTGGGGTCCCATCANGGTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCAT
CAGCAGCCTGCAACCTGNGGATTTTGCAACTTACTACTGTCAACAGAGTTACAGTAACCC]T[CTC
ACTTTCGGCGGNGGGACCAANGTGGAGATCAAAC]GAACTGTGGCTGCACCATCTGTCTTCATCT
TCCCGCCATCTGATGAGCAGTTGAAATCTGGAACTGCCTCTGTTGTGTGCCTGCTGAATAACTTCT
ATCCCAGAGAGGGCCAAAGTACA

FIG. 16H



International application No PCT/US96/05928

			
1	ASSIFICATION OF SUBJECT MATTER		
IPC(6)	.A61K 39/00, C07K 16/18, C12N 5/16, 15/13, :424/184,1; 514/12; 536/23,53; 530/387,1.		
	to International Patent Classification (IPC) or to be	th national classification and IPC	
B. FIE	LDS SEARCHED		
Minimum e	locumentation searched (classification system follow	red by classification symbols)	
U.S.	424/184.1; 514/12; 536/23.53; 530/387.1.		
Documenta	tion searched other than minimum documentation to	the extent that such documents are include	d in the fields searched
Electronic o	lata base consulted during the international search (name of data base and, where practicable	:, search terms used)
APS, DIA	ALOG		
C. DOC	UMENTS CONSIDERED TO BE RELEVANT		
Category®	Cuation of document, with indication, where	appropriate, of the relevant passages	Relevant to claim No.
X	EP 463151 A; (JAKOBOVITZ ET , abstract.	AL) 01 February 1991, See	1-4,19
Υ			1-44
Y,X	Expert Opinion on Investigational	Drugs, Volume 3, Number	20-44
	3, issued March 1994, Emery et a	I., "Humanised monoclonal	20 44
	antibodies for therapeutic applic	cations", pages 241-251,	
	especially pages 243-245.	, , , , , , , , , , , , , , , , , , , ,	•
		}	
İ			
		ļ	
1			
Facus	et documents are insect in the continuation of Boy (See patent family annex	
	attender of the state of the st	tace document published after the inte	
v	municipal and the territor has a discord which is now supplied at	data and not in contlict with the advisor approxi-	continue and or matershap the
	ice document processed on or acts office advantage of thing state	"X" observation of particular relevance, the considered novel or cannot be consider	claimed invention cannot be ed to involve in inventive step
- 1:	orient which may throw doubts on pounds channed to which is I to establish his problemation date of another estation or other of travels as producing	when the document is taken alone	
•	mischer er betreng to an out dear excession of the exceptional and court	V document of particular relevance, the considered to involve an inventor confined with one or more other so them; one one necessor skided in the	step when the document is extrementation
	forming alrested to a great of the entire and the same conclusive man- prior to the entire and	"A succument mendicr of the same paten-	tatusts
has in the .	ectual compilation of are interferomant serior.	Date of mading of the international sea	reh report
: " (1 , Y)	99(1	09 AUG1996	
	and the forms and the control	Substitute 1 //3/07/10	7-30/10
des PCT Washington	Bx 1/25.	THE A PLISE.	
comme No		Terestorie No. 1923 (1980) 196	
PC . 48	A large second manager, by a religious		